

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
25 March 2004 (25.03.2004)

PCT

(10) International Publication Number  
**WO 2004/023891 A1**

(51) International Patent Classification<sup>7</sup>: **A23L 1/054**,  
1/308

(21) International Application Number:  
PCT/EP2003/009949

(22) International Filing Date:  
8 September 2003 (08.09.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0221184.5 13 September 2002 (13.09.2002) GB

(71) Applicant (*for all designated States except US*): CER-  
ESTAR HOLDING B.V. [NL/NL]; Nijverheidsstraat 1,  
P.O. Box 9, NL-4551 La Sas van Gent (NL).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): VERCAUTEREN,  
Ronny, Leontina, Marcel [BE/BE]; Bergstraat 29, B-9120  
Beveren (BE). NGUYEN, Van Sau [BE/BE]; Rue Guil-  
laume Lekeu 46, B-1070 Brussels (BE).

(74) Agent: WILKINSON, Stephen, John; Stevens, Hewlett  
& Perkins, 1 St. Augustine's Place, Bristol BS1 4UD (GB).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,  
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


**Declaration under Rule 4.17:**

— *of inventorship (Rule 4.17(iv)) for US only*

**Published:**

— *with international search report*

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*



WO 2004/023891 A1

(54) Title: SLOWLY DIGESTIBLE CARBOHYDRATE MATERIALS FOR USE IN FOOD AND DRINK COMPOSITIONS

(57) Abstract: Use of a carbohydrate material comprising a glycosidic backbone of which at least 5% consists of  $\alpha$ -1,6-D-glucopyranosidic linkages as a food and/or drink ingredient which, when subjected to hydrolysis in the presence of intestinal enzymes, undergoes hydrolysis at a rate which is not greater than 0.8 times the hydrolysis rate of maltose under the same conditions such that low levels of glucose are released over prolonged periods of time. This carbohydrate material is incorporated in food and drink compositions at 15% or more by weight based on the total weight of the composition.

**SLOWLY DIGESTIBLE CARBOHYDRATE MATERIALS**  
**FOR USE IN FOOD AND DRINK COMPOSITIONS**

The present invention relates to the use of certain specified carbohydrate materials in food and drink compositions and to food and drink compositions containing the specified carbohydrate materials.

It is common knowledge that oligomers of glucose that are aligned to each other by an  $\alpha$ -1,4- glucosidic bond are easily hydrolysed and digested in the first parts of the digestion channel (mouth-stomach-small intestine) of humans and animals. Classical examples of such compounds are the commercially available hydrolysis products of starch, for example maltose and maltodextrins. These compounds give a rapid increase in glucose blood level after oral ingestion similar to glucose itself. Maltose is a disaccharide composed of two anhydroglucopyranoside units connected to each other by an O- $\alpha$ -1,4- linkage.

Maltodextrins are hydrolysis products of starch having a dextrose equivalent (D.E.) lower than 20. The term dextrose equivalent indicates the percentage of reducing sugars, expressed as glucose on dry basis.

U.S. Patent 2,983,613 describes the addition of dextran to bread doughs to improve the softness and shelf-life of the baked bread. The dextran may, according to this document, be used in doughs in amounts equal to from about 0.01 to about 10% by weight, based on the weight of the flour content of the dough, although it is stated that there is no advantage in employing more than 5% by weight.

The specified carbohydrate material is defined as a carbohydrate material which provides a slower release and, therefore, a slower absorption into the body, of glucose during the transit through the small intestine as compared to maltose.

Applications of the specified carbohydrate material are manifold, i.e. they can be used to deliver to the diabetic patient sufficient carbohydrates without a significant raise of the serum glucose level. Also, the addition of the specified carbohydrate material to the diet of elderly people confronted with a reduced glucose tolerance could have a beneficial effect.

In the domain of sports nutrition, the specified carbohydrate material could have interesting applications. The carbohydrate material could supply an athlete with a steady and constant carbohydrate supply during physical exercise. It could also be used in foods and/or supplements for growing children and in so called "energy drinks".

The present invention is based on the discovery that a certain amount of  $\alpha$ -1,6-D-glucopyranosidic linkages are to be used as the carbohydrate materials. According to the invention, the carbohydrate material comprises a glycosidic backbone of which at least 5% consists of  $\alpha$ -1,6-D-glucopyranosidic linkages. Preferably, at least 10% of the glycosidic backbone in the carbohydrate material consists of  $\alpha$ -1,6-D-glucopyranosidic linkages. In a more preferred embodiment, at least 30% of the glycosidic backbone in the carbohydrate material consists of  $\alpha$ -1,6-D-glucopyranosidic linkages.

According to a first aspect, the invention provides the use of a carbohydrate material comprising a glycosidic backbone of which at least 5% consists of  $\alpha$ -1,6-D-glucopyranosidic linkages as a food and/or drink ingredient which, when subjected to hydrolysis in the presence of intestinal enzymes, undergoes hydrolysis at a rate which is not greater than 0.8 times the hydrolysis rate of maltose under the same conditions such that low levels of glucose are released over prolonged periods of time.

More preferably, the carbohydrate material, when subjected to hydrolysis in the presence of intestinal enzymes for a period of time not exceeding 6 hours undergoes

hydrolysis at a rate which is in the range of from 0.1 to 0.6 times the hydrolysis rate of maltose under the same hydrolysing conditions.

The carbohydrate material preferably comprises at least one oligosaccharide or polysaccharide having a molecular weight in the range of from 300 to 1,000,000 daltons. The carbohydrate material may comprise one oligosaccharide or polysaccharide or may comprise a mixture of two or more oligosaccharides or polysaccharides. As stated above, the carbohydrate material comprises a glycosidic backbone of which at least 5% consists of  $\alpha$ -1,6-D-glucopyranosidic linkages.

In the case where the carbohydrate material comprises a mixture of two or more different oligosaccharides or polysaccharides, 5% of the total glycosidic backbone components in the mixture should comprise  $\alpha$ -1,6-D-glucopyranosidic linkages. According to a preferred embodiment of the invention, the carbohydrate material is selected from isomaltose, dextrans, pullulans, alternans and mixtures of two or more of these. According to a more preferred embodiment, the carbohydrate material comprises a mixture of isomaltose and one or more of a dextran, pullulan or alternan.

A dextran is a polyglucose and dextrans of a wide variety of molecular weights and structures have been known for many years. The dextran produced by *Leuconostoc mesenteroides* NRRL B-512F has about 95% of  $\alpha$ -1,6- linkages, the other 5% being  $\alpha$ -1,3- linkages and make up the branches. Examples of dextrans that may be used according to the invention include synthetic dextrans and native dextrans such as those produced by *Leuconostoc mesenteroides* NRRL B-742, one with 14%  $\alpha$ -1,4- branch linkages and 86%  $\alpha$ -1,6- linkages and a dextran with 50%  $\alpha$ -1,3- branch linkages and the remainder being  $\alpha$ -1,6- linkages. Other examples are dextrans produced by *L. mesenteroides* NRRL B-1355. This *Leuconostoc* strain produces two polysaccharide types: a dextran with 95%  $\alpha$ -1,6- linkages in the main chain and 5%  $\alpha$ -1,3- branched linkages and another composed of alternating  $\alpha$ -1,6- linkages and  $\alpha$ -1,3- linkages in the main glycosidic chain and has 11%  $\alpha$ -1,3- branched linkages. Another example is the water soluble dextran produced by

*Streptococcus mutans* 6715 which is composed of 64%  $\alpha$ -1,6- linkages in the main glycosidic chain and 36%  $\alpha$ -1,3- branched linkages. Yet another example is the dextran produced by *Gluconobacter oxidans* ATCC 11894 which has 90%  $\alpha$ -1,6- linkages, the remaining being  $\alpha$ -1,4- linkages.

The carbohydrate material may comprise a pullulan. Pullulan is a viscous polysaccharide that may be extracellularly produced by growing certain yeasts on sugars and comprises a polymer of D-glucopyranose residues, containing  $\alpha$ -1,4- linkages and  $\alpha$ -1,6- linkages in the ratio 2:1. The structure is that of a polymer of maltotriose, connected end-to-end by an  $\alpha$ -1,6- glycosidic linkage. In addition to maltotriose units, pullulan also has 5 to 7% maltotetraose units distributed along the main chain. Thus, about 33% of the glycosidic linkages in pullulan are  $\alpha$ -1,6- linkages, the remainder being  $\alpha$ -1,4- linkages.

Alternatively, the carbohydrate material may be, or comprise, an alternan. Alternan is a homopolymer of glucose and has alternating  $\alpha$ -1,6- and  $\alpha$ -1,3- linked glucose residues in the main chain with varying amounts of  $\alpha$ -1,3- branch linkages.

The present invention also provides a food or drink composition which contains as one of its components the carbohydrate material as described above in an amount greater than 15% by weight based on the total weight of the composition. Typically, the food or drink composition may be a diabetic food or drink, a baby food or drink, a diet food or drink, for instance for sedentary people, a specially formulated food or for people having a reduced glucose tolerance, for instance the elderly, or a food or drink for growing children. The carbohydrate material may, advantageously, be used as a component in sports or energy beverages. Typically, the carbohydrate material will be present in a composition in an amount of at least 20% by weight.

The presence of the carbohydrate material in food and drink compositions provides, as mentioned earlier, a slower release, and therefore a slower absorption into the body, of glucose. This is correlated with a low glycemic index. Accordingly,

the present invention further provides the use of a carbohydrate material comprising a glycosidic backbone of which at least 5%, preferably at least 10% and more preferably at least 30%, consists of  $\alpha$ -1,6-D-glucopyranosidic linkages in the manufacture of food and drink compositions having a low glycemic index. It follows that it is possible to reduce the glycemic index of a food or drink composition by replacing at least part of the carbohydrate content thereof with a slowly digestible carbohydrate material comprising a glycosidic backbone of which at least 5% consists of  $\alpha$ -1,6-D-glucopyranosidic linkages, as disclosed herein before.

**Example 1: In-vitro hydrolysis of maltose and isomaltose by rat intestinal acetone powder**

A 1% substrate solution (w/w) of maltose or isomaltose was prepared in 0.05 M phosphate buffer pH 6 and equilibrated at 37°C for 10 minutes. A suspension of 2.5% rat intestinal acetone powder (Sigma) was used as a source of glucoamylase-maltase complex and sucrase-isomaltase and was prepared in 0.05M phosphate buffer pH 6 and equilibrated at 37°C for 10 minutes. 0.6ml rat intestinal acetone powder (excess) was added to 6 ml of substrate solution and mixed. The mixture was incubated at 37°C and a 1.5 ml sample was taken (0 hours incubation time). Further samples were taken after 2, 4 and 6 hours of incubation. The samples were diluted with 4 ml of demineralised water and boiled for 5 minutes. After the denaturation step, each sample was filtered through a 0.45  $\mu$ m filter.

The filtrate was sent through a Dionex OnGuard-ATM filter. Glucose was determined by HPLC. The results for the in-vitro small intestinal digestion tests for maltose and isomaltose are given in Table 1.

**Table 1****In vitro hydrolysis of maltose and isomaltose by rat intestinal acetone powder**

Compound	% glucose produced		
	2 Hours	4 Hours	6 Hours
Maltose	100.0	100.0	100.0
Isomaltose	49.0	52.0	58.0

This example shows the hydrolysis kinetics of isomaltose. This example shows a slower release of glucose compared to maltose.

**Example 2: In-vitro hydrolysis of dextrans by rat intestinal acetone powder**

The same experimental procedure as in Example 1 was followed, except that the 2.5% rat intestinal acetone powder solution was replaced by a 10% rat intestinal acetone powder solution. Dextrans of different molecular weight were used as substrates. Dextrans with Mw ~1000, ~5000, ~12000, ~25000, ~50000, ~80000, ~150000 and ~270000 were from Fluka. The dextrans T-500 (Mw~500000) and T-2000 (Mw~2000000) were from Amersham Pharmacia Biotech. The results are presented in Table 2.

**Table 2**

Compound	% glucose produced		
	2 Hours	4 Hours	6 Hours
Maltose	100.0	100.0	100.0
Isomaltose	61.1	75.5	84.5
Dextran 1000	38.9	54.0	64.4
Dextran 5000	34.3	46.7	55.0
Dextran 12000	24.8	33.5	39.5
Dextran 25000	18.6	25.4	30.1
Dextran 50000	15.4	20.8	24.9
Dextran 80000	14.5	19.0	22.3
Dextran 150000	12.3	17.2	20.6
Dextran 270000	11.8	16.9	20.2
Dextran T 500 Pharmacia	12.7	17.9	20.3
Dextran T2000 Pharmacia	11.8	16.3	19.0

This example shows the hydrolysis kinetics of a series of dextran polymers having different degrees of polymerisation. This example shows a slower release of glucose compared to maltose.

**Example 3: In-vitro hydrolysis of pullulans by rat intestinal acetone powder**

The same experimental procedure as in Example 2 was followed. Pullulans with Mw ~5000, ~10000, ~20000, ~50000, ~100000, ~200000, ~400000, ~800000 from Shodex and isomaltose were used as substrates. The results are presented in Table 3.

**Table 3**

Compound	% glucose produced		
	2 Hours	4 Hours	6 Hours
Maltose	100.0	100.0	100.0
Isomaltose	65.7	79.1	87.5
P-5	44.2	57.5	70.4
P-10	26.9	37.1	46.0
P-20	18.8	26.0	32.8
P-50	13.3	18.6	23.3
P-100	10.1	14.6	18.5
P-200	8.6	12.6	16.6
P-400	8.0	11.7	15.1
P-800	7.7	11.5	14.6

This example shows the hydrolysis kinetics of a series of pullulan polymers having different degrees of polymerisation. The example shows a slower release of glucose compared to maltose.



**Example 4: Sport or isotonic drink composition****Ingredients for one litre of drink:**

Carbohydrate Material*	30g
Sucrose	40g
Sodium chloride	1.27g
Citric acid	1g
Orange flavour	2g
Sodium benzoate (10%w/v)	1.5ml
Water	to one litre

---

\* Any carbohydrate material comprising a glycosidic backbone of which at least 5% comprises a  $\alpha$ -1,6-D-glucopyranosidic linkages.

For the preparation of the drink, the sucrose and the carbohydrate material were dissolved into water and then all other ingredients were added to the solution obtained.

**Example 5: Cereal bar formulation and method of manufacture**

<u>Ingredients</u>	<u>%</u>
Cereal flakes	30
Carbohydrate material	20
Rice crispies	8
Raisins	5
Binding syrup (see below)	37

<u>Binding syrup formulation</u>	<u>%</u>
Glucose syrup (9% fructose)	27.9
Sucrose	15.2
Maltodextrin (6DE)	13.7
Invert sugar	12.9
Glycerol	10.8
Margarine	7.8

Sorbitol syrup	3
Lecithin	2.6
Water	4.8
Salt	1.3
Vanilla	as needed

---

\* Any carbohydrate material comprising a glycosidic backbone of which at least 5% comprises a  $\alpha$ -1,6-D-glucopyranosidic linkages.

#### Manufacturing process of cereal bars

The ingredients of the binding syrup were blended and to this blend was added the carbohydrate material. The mixture was then heated to 100°C. A blend of the remaining ingredients (cereal flakes, rice crispies and raisins) was prepared. The binding syrup was then added and mixed with the other ingredients until a uniform mixture was obtained. The mixture was then formed into bars and packaged.

## CLAIMS

1. Use of a carbohydrate material comprising a glycosidic backbone of which at least 5% consists of  $\alpha$ -1,6-D-glucopyranosidic linkages as a food and/or drink ingredient which, when subjected to hydrolysis in the presence of intestinal enzymes, undergoes hydrolysis at a rate which is not greater than 0.8 times the hydrolysis rate of maltose under the same conditions such that low levels of glucose are released over prolonged periods of time.
2. Use according to claim 1, characterised in that the carbohydrate material, when subjected to hydrolysis in the presence of intestinal enzymes for not greater than 6 hours, undergoes hydrolysis at a rate which is in the range of from 0.1 to 0.6 times the hydrolysis rate of maltose under the same conditions.
3. Use according to either claim 1 or claim 2, characterised in that the intestinal enzymes are selected from glucoamylase-maltase, sucrase-isomaltase and mixtures thereof.
4. Use according to any one of claims 1 to 3, characterised in that the carbohydrate material comprises a glycosidic backbone of which at least 10% consists of  $\alpha$ -1,6-D-glucopyranosidic linkages.
5. Use according to claim 4, characterised in that the carbohydrate material comprises a glycosidic backbone of which at least 30% consists of  $\alpha$ -1,6-D-glucopyranosidic linkages.
6. Use according to any one of claims 1 to 5, characterised in that the carbohydrate material comprises at least one oligosaccharide, polysaccharide or mixtures thereof having a molecular weight in the range of from 300 to 1,000,000 daltons.

7. Use according to claim 6, characterised in that the carbohydrate material is selected from isomaltose, dextrans, pullulans, alternans and mixtures of two or more of these.
8. Use according to any one of claims 1 to 7, as an ingredient for diabetic foods, baby foods, special dietary formulations and/or sports drinks.
9. Use according to any one of claims 1 to 8, characterised in that the carbohydrate material is added in amounts of at least 15% by weight to said food and/or drink compositions.
10. Use according to claim 9, characterised in that the carbohydrate material is added in amounts of at least 20% by weight to said food and/or drink compositions.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 03/09949

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 A23L1/054 A23L1/308

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A23L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

PAJ, WPI Data, EPO-Internal, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 153 013 A (FISONS PLC) 28 August 1985 (1985-08-28) claims; examples	1-10
X	US 5 116 820 A (HIJI YASUTAKE) 26 May 1992 (1992-05-26) column 1, line 32 - line 46; claims column 5, line 26 - column 6, line 12	1-8
X	US 4 629 725 A (HIJI YASUTAKE) 16 December 1986 (1986-12-16) the whole document	1-8
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

19 December 2003

Date of mailing of the international search report

02/01/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Lepretre, F

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 03/09949

G.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DATABASE WPI Section Ch, Week 198432 Derwent Publications Ltd., London, GB; Class D13, AN 1984-198539 XP002234400 &amp; JP 59 113856 A (ENDO A), 30 June 1984 (1984-06-30) abstract</p>	1
A	<p>WO 00 70964 A (BAUCHE ANNE ;NESTLE SA (CH); SCHMID DANIEL (CH); ARRIGONI EVA (CH)) 30 November 2000 (2000-11-30) claims</p>	1-10
A	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1997 GRIMBLE GEORGE K ET AL: "Differences in the glycaemic response to dextran and maltodextrin ingestion in man." Database accession no. PREV199799727067 XP002234399 abstract &amp; PROCEEDINGS OF THE NUTRITION SOCIETY, vol. 56, no. 2, 1997, page 225A Joint Meeting of the Clinical Nutrition and Metabolism Group of the Nutrition Society and the British Association for Parenteral and Enteral Nutrition;Blackpool, England, UK; December 3-5, 1996 ISSN: 0029-6651</p>	1
A	<p>GB 830 951 A (COMMW ENGINEERING COMPANY) 23 March 1960 (1960-03-23)</p>	

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/09949

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0153013	A	28-08-1985	AT 56148 T	15-09-1990
			AU 571641 B2	21-04-1988
			AU 3819385 A	08-08-1985
			DE 3579471 D1	11-10-1990
			DK 39685 A	02-08-1985
			EP 0153013 A2	28-08-1985
			JP 60188403 A	25-09-1985
US 5116820	A	26-05-1992	JP 1642389 C	28-02-1992
			JP 3008188 B	05-02-1991
			JP 62236469 A	16-10-1987
US 4629725	A	16-12-1986	JP 59074976 A	27-04-1984
			JP 61053029 B	15-11-1986
			US 4913925 A	03-04-1990
JP 59113856	A	30-06-1984	NONE	
WO 0070964	A	30-11-2000	EP 1060673 A1	20-12-2000
			AU 5968900 A	12-12-2000
			CA 2374041 A1	30-11-2000
			WO 0070964 A1	30-11-2000
			NZ 515196 A	26-09-2003
			ZA 200109013 A	31-01-2003
GB 830951	A	23-03-1960	NONE	